

Patient:	
Full name:	Nationality:
Address:	Date of birth:
Zip code/city:	Male Female
Country:	
The Patient mentioned above	is being treated for:
Rheumatoid arthritis	Non-radiographic axial spondyloarthritis
Juvenile idiopathic arthritis	Psoriasis
Psoriatic arthritis	Juvenile psoriasis
Ankylosing spondylitis	Other:
Is treated by:	
Rheumatologist / Dermatologist:	
Hospital:	
Patient uses: Etanercept (Enbrel®) 50 mg subcutaneously (sc) once weekly Etanercept (Enbrel®) 50 mg sc twice weekly Etanercept (Enbrel®) 25 mg sc twice weekly Etanercept (Enbrel®) — mg sc once / twice weekly	25 og Presided Sinder En bre i 25 og Charactery 10 The standard Sinder En bre i 25 og Charactery 10 The standard Sinder En bre i 25 og Charactery 10 The standard Sinder En bre i 25 og Charactery 10 The standard Sinder En bre i 25 og Charactery 10 The standard Sinder The standard Sindard Sind
The patient self-administers the medication. Therefore, this syringes, or packs with syringes, needles and vials with Enbre The medication needs to be stored between 2°C and 8°C. Thing a coolpack with cooling liquid. Alternatively, the medicating is period of up to 4 weeks, after which it should not be reused within 4 weeks after removal from refrigeration.	el powder (see picture above) in his/her hand luggage. Herefore, the medication is transported in a coolbag, contai-
(Doctor's signature)	(Doctor's stamp)